

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 978 512 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
09.02.2000 Bulletin 2000/06

(21) Application number: 98401944.8

(22) Date of filing: 29.07.1998

(51) Int. Cl.⁷: **C07D 295/088**, C07C 211/08,
C07D 211/04, C07D 295/185,
C07D 211/62, C07D 211/70,
C07D 207/20, A61K 31/13,
A61K 31/40, A61K 31/445,
A61K 31/495

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(71) Applicant:
SOCIETE CIVILE BIOPROJET
F-75003 Paris (FR)


(72) Inventors:
• **Schwartz, Jean-Charles**
75014 Paris (FR)
• **Arrang, Jean-Michel**
91419 Dourdan (FR)
• **Garbarg, Monique**
75017 Paris (FR)
• **Lecomte, Jeanne-Marie**
75003 Paris (FR)

- **Ligneau, Xavier**
75013 Paris (FR)
- **Schunack, Walter G.**
14129 Berlin (DE)
- **Stark, Holger**
12157 Berlin (DE)
- **Ganellin, Charon Robin**
Welwyn, Hert AL6 OTD (GB)
- **Leurquin, Fabien**
London E2 6DZ (GB)
- **Sigurd, Elz**
12107 Berlin (DE)

(74) Representative:
Le Guen, Gérard et al
CABINET LAVOIX
2, place d'Estienne d'Orves
75441 Paris Cédex 09 (FR)

(54) **Non-imidazole aryloxy (or arylthio) alkylamines as histamine H₃-receptor antagonists and their therapeutic applications**

(57) Compounds of formula (I):



and their use for preparing medicaments acting as antagonists at the H₃-receptors of histamine.

EP 0 978 512 A1

Description

[0001] The present invention relates to novel aryloxy (or arylthio) alkylamines, to their preparation and to their therapeutic applications.

[0002] Antagonists of histamine H_3 receptor are known especially to increase synthesis and release of cerebral histamine. Through this mechanism, they induce an extended wakefulness, an improvement in cognitive processes, a reduction in food intake and a normalization of vestibular reflexes (Schwartz et al., *Physiol. Rev.*, 1991, 71; 1-51).

[0003] Whence these agents are potentially useful in several central nervous system disorders such as Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.

[0004] All the H_3 receptor antagonist compounds known so far resemble histamine in possessing an imidazole ring (Ganellin et al., *Ars Pharmaceutica*, 1995, 36:3, 455-468; Stark et al., *Drug of the Future*, 1996, 21(5), 507-520).


[0005] Nevertheless, such imidazole derivatives may show drawbacks such as poor blood-brain barrier penetration and/or some hepatic and ocular toxicities. These drawbacks, which can prevent their therapeutic development, appear to be linked to the presence of an imidazole ring substituted by a hydrophobic chain.

[0006] Attempts to develop H_3 receptor antagonists without an imidazole ring have up to now been unsuccessful, the compounds being of low potency.

[0007] In this respect, non-imidazole compounds such as betahistine (J-M. Arrang et al., *Eur. J. Pharmacol.* 1985, 111: 72-84), phencyclidine (J-M. Arrang et al., *Eur. J. Pharmacol.* 1988, 157: 31-35), dimaprit (J-C Schwartz et al., *Agents Actions* 1990, 30: 13-23), clozapine (M. Kathmann et al., *Psychopharmacology* 1994, 116: 464-468), and sesquiterpenes (M. Takigawa et al., JP 06 345 642 (20 Dec 1994)) were suggested to display H_3 -receptor antagonism but all these compounds have only very low potency.


[0008] The present invention provides new compounds, the structure of which do not contain an imidazole moiety, which are useful as histamine H_3 -receptor antagonists while avoiding the above-mentioned drawbacks of the known H_3 -antagonists.

[0009] The compounds of the invention have the following general formula (I):




in which:

- C_nH_{2n} is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- R^1 and R^2 may be identical or different and represent each independently
 - a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
 - a saturated nitrogen-containing ring



with m ranging from 4 to 7, or

- an unsaturated nitrogen-containing ring



with p, q and r being 1 to 3 independently, such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or

- a morpholino group, or
- a N-substituted piperazino group:



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- n_3 is an integer from 0 to 5; ← An integer? not a atom or chain?
- R^3 represents each independently

- a halogen atom,
- a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino, α -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
- or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring.

[0010] The invention also relates to the addition salts which the compounds form with pharmaceutically acceptable acids. The pharmaceutically acceptable salts comprise the nontoxic salt of inorganic or organic acids. Examples of these salts include the hydrochloride, the hydrobromide or the hydrogen maleate or hydrogen oxalate.

[0011] The present invention also encompasses the hydrates of the compounds, the hydrated salts of these compounds and the polymorphic crystalline structures. When the compounds can exist in one or a number of isomeric forms according to the number of asymmetric centres in the molecule, the invention relates both to all the optical isomers and to their racemic modifications and the corresponding diastereoisomers. The separation of the diastereoisomers and/or of the optical isomers can be carried out according to methods known per se.

[0012] According to the invention, lower alkyl or cycloalkyl is intended to mean a linear or branched alkyl group containing from 1 to 6 carbon atoms, or a saturated carbocycle containing 3 to 6 carbon atoms.

[0013] Typically examples of lower alkyl are methyl, ethyl, propyl, isopropyl and butyl groups.

[0014] A preferred group of compounds according to the invention comprises those with R^1 and R^2 representing independently a lower alkyl group, especially an ethyl group.


[0015] Preferred compounds are also those of formula (I) in which R^1 and R^2 taken together with the nitrogen atom to which they are attached, form a saturated nitrogen-containing ring:



especially with m being 4, 5 or 6, optionally substituted with an alkyl group, preferably a methyl group.

[0016] Another preferred group of compounds comprises compounds (I) in which R^1 and R^2 taken together with the nitrogen atom to which they are attached, form an unsaturated nitrogen-containing ring:

ii)



especially with p, q, and r being 1 or 2. In this group, more preferred compounds are those with p being 2 and q and r each being 1.

[0017] Typical example of $-NR^1R^2$ representing a N-substituted piperazino group is N-acetyl/piperazino.

[0018] A preferred group of compounds according to the invention is the group composed of compounds of formula (I) in which X is an oxygen atom.

[0019] Another preferred group of compounds comprises compounds (I) in which $-C_nH_{2n}-$ is a linear chain $-(CH_2)_n-$ with n being as previously defined.

[0020] Preferred compounds are also those with n varying from 3 to 5, and with n being more preferably 3.

[0021] A sub-class of compounds according to the invention comprises the compounds of formula (I) with n_3 being zero that is those having an unsubstituted phenyl moiety.

[0022] Another group of compounds according to the invention is composed of compounds containing one or more substituents R^3 which may be identical or different. In this group, the compounds having a mono- or di-substituted ($n_3 = 1$ or 2) phenyl moiety are preferred and those mono-substituted with one group R^3 as defined above in para-position are particularly preferred.

[0023] Among these compounds, (n_3 being 1) R^3 is preferably a halogen atom or a cyano, nitro, alkanoyl, alkyloximino or α -hydroxyalkyl group.

[0024] Still more preferred compounds are those with R^3 being CN, NO_2 , $COCH_3$, COC_2H_5 , $H_3C-C=N-OH$, $H_3C-CH-OH$.

[0025] R^3 being a halogen atom may be advantageously selected from fluorine, chlorine and bromine.

[0026] R^3 being an aryl group, may be especially a phenyl group.

[0027] In the other substituents R^3 , the aryl moiety is advantageously a phenyl moiety.

[0028] R^3 being an aryloxy group may be especially a phenoxy group.

[0029] According to the invention, alkanoyl is intended to mean a group containing an alkyl moiety as defined above.

[0030] Typical examples of R^3 being an alkanoyl, aroyl or aryalkanoyl group are acetyl, butyryl and propionyl groups, benzoyl group or phenylacetyl group.

[0031] Typical examples of R^3 forming together with the carbon atoms of the phenyl ring to which it is fused, a saturated ring leads to 5,6,7,8-tetrahydronaphthyl or forming a benzene ring leads to a naphthyl moiety.

[0032] According to the invention, alkenyl or alkynyl group may contain advantageously from 1 to 8 carbon atoms, in particular from 1 to 6 carbon atoms and preferably 1 to 4 carbon atoms.

[0033] In carboalkoxy, carboxyamido or carboxamide groups, the hydrocarbon chain is saturated, linear or branched and contains an alkyl moiety as defined above.

[0034] In alkoxy, alkyloximino, arylalkyl or α -hydroxyalkyl group, the alkyl moiety is as previously defined also.

[0035] Particularly preferred compounds are:

- 1-(5-phenoxy-pentyl)-piperidine
- 1-(5-phenoxy-pentyl)-pyrrolidine
- N-methyl-N-(5-phenoxy-pentyl)-ethylamine
- 1-(5-phenoxy-pentyl)-morpholine
- N-(5-phenoxy-pentyl)-hexamethyleneimine
- N-ethyl-N-(5-phenoxy-pentyl)-propylamine
- 1-(5-phenoxy-pentyl)-2-methyl-piperidine
- 1-(5-phenoxy-pentyl)-4-propyl-piperidine
- 1-(5-phenoxy-pentyl)-4-methyl-piperidine
- 1-(5-phenoxy-pentyl)-3-methyl-piperidine
- 1-acetyl-4-(5-phenoxy-pentyl)-piperazine
- 1-(5-phenoxy-pentyl)-3,5-trans-dimethyl-piperidine
- 1-(5-phenoxy-pentyl)-3,5-cis-dimethyl-piperidine

- 1-(5-phenoxy-pentyl)-2,6-cis-dimethyl-piperidine
 4-carboethoxy-1-(5-phenoxy-pentyl)-piperidine
 3-carboethoxy-1-(5-phenoxy-pentyl)-piperidine
 1-(5-phenoxy-pentyl)-1,2,3,6-tetrahydropyridine
 5 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
 10 1-[5-(2-naphthyl-oxy)-pentyl]-pyrrolidine
 1-[5-(1-naphthyl-oxy)-pentyl]-pyrrolidine
 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
 1-[5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
 15 1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
 1-(5-phenoxy-pentyl)-2,5-dihydropyrrole
 1-[5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
 1-(4-phenoxybutyl)-pyrrolidine
 1-(6-phenoxyhexyl)-pyrrolidine
 20 1-(5-phenylthiopentyl)-pyrrolidine
 1-(4-phenylthiobutyl)-pyrrolidine
 1-(3-phenoxypropyl)-pyrrolidine
 1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
 25 1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
 1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
 1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
 N-[3-(4-nitrophenoxy)-propyl]-diethylamine
 30 N-[3-(4-cyanophenoxy)-propyl]-diethylamine
 1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
 1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine
 N-[3-(4-acetylphenoxy)-propyl]-diethylamine
 1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
 35 1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
 1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine
 1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
 1-[5-(4-cyanophenoxy)-pentyl]-piperidine
 40 N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
 N-[2-(4-cyanophenoxy)-ethyl]-diethylamine
 N-[3-(4-cyanophenoxy)-propyl]-dimethylamine
 N-[4-(4-cyanophenoxy)-butyl]-diethylamine
 N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
 45 1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
 N-[6-(4-cyanophenoxy)-hexyl]-diethylamine
 N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
 50 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine
 55 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
 1-[3-(4-formylphenoxy)-propyl]-piperidine

1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
 N-[3-(4-propionylphenoxy)-propyl]-diethylamine
 1-[3-(4-butyrylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine

5

[0036] More preferred compounds are:

1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 N-[3-(4-cyanophenoxy)-propyl]-diethylamine
 10 N-[3-(4-acetylphenoxy)-propyl]-diethylamine
 1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine
 N-[4-(4-cyanophenoxy)-butyl]-diethylamine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
 15 N-[3-[4-(1-hydroxyethyl)-phenoxy]-propyl]-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine

20

[0037] Compounds of formula (I) in which:

- -NR¹R² is a pyrrolidinyl group, C_nH_{2n} is a linear chain -(CH₂)_n- and n₃ is zero, X being an oxygen atom with n ranging from 3 to 5, or X being a sulfur atom with n being 4 or 5;
- 25 • -NR¹R² is a piperidinyl group, C_nH_{2n} is a linear chain -(CH₂)_n- and X is an oxygen atom, n₃ being zero with n being 2, 5 or 8 or n₃ being 1 with R³ being 4-CN and n being 5;
- -NR¹R² is a diethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain -(CH₂)_n- and n₃ is 1, R³ being 4-NO₂ or 4-COCH₃ with n being 3 or R³ being 4-CN with n being 2 to 4;
- 30 • -NR¹R² is a dimethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain -(CH₂)_n- and n₃ is 1, R³ being 4-CN with n being 3,

are known in the art.

[0038] A subject of the invention is thus the use of these compounds as antagonists at the histamine H₃-receptors, in particular to prepare medicaments acting as H₃-antagonists intended for the treatments detailed below.

35 [0039] The compounds according to the invention may be prepared according to one of the following schemes 1-5:


40

45


50

55


SCHEME 1 (methods A, B, C, D, H and K):




SCHEME 2 (methods F and L):




SCHEME 3 (method E):



SCHEME 4 (method G):



SCHEME 5 (method J):



[0040] In these schemes, R¹, R², R³, X and n are as defined in general formula (I).

[0041] Me and Et are intended to mean methyl and ethyl.

[0042] Detailed synthesis procedures are given in the examples.

[0043] The compounds of formula (I) according to the invention have antagonistic properties at the histamine H₃-receptors. They cause an increase in synthesis and release of cerebral histamine.

[0044] This property makes the compounds of the invention useful derivatives in human or veterinary medicine.

5 [0045] Their therapeutical applications are those known for H₃-antagonist compounds and especially relate to the central nervous system disorders such as Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.

[0046] Therefore, the compounds of formula (I) according to the invention are advantageously used as active ingredient of medicaments which act as an antagonist of H₃-receptors of histamine, in particular of medicaments having psychotropic effects, promoting wakefulness, attention, memory and improving mood, in treatment of pathologies such as 10 Alzheimer disease and other cognitive disorders in aged persons, depressive or simply asthenic states.

[0047] Their nootropic effects can be useful to stimulate attention and memorization capacity in healthy humans.

[0048] In addition, these agents can be useful in treatment of obesity, vertigo and motion sickness.

[0049] It can also be useful to associate the compounds of the invention with other psychiatric agents such as neuroleptics to increase their efficiency and reduce their side effects.

15 [0050] Application in certain form of epilepsy is also foreseen.

[0051] Their therapeutic applications involve also peripheral organs mainly a stimulant of secretions or gastro-intestinal motricity.

[0052] The compounds of the invention are particularly useful for the treatment of CNS disorders of aged persons.

20 [0053] The present invention also relates to medicaments having the above-mentioned effects comprising as active ingredient, a therapeutically effective amount of a compound of formula (I).

[0054] The present invention also relates to pharmaceutical compositions containing as active ingredient, a therapeutically effective amount of a compound (I) together with a pharmaceutically acceptable vehicle or excipient.

[0055] The medicaments or pharmaceutical compositions according to the invention can be administered via oral, parenteral or topical routes, the active ingredient being combined with a therapeutically suitable excipient or vehicle.

25 [0056] According to the invention, oral administration is advantageously used.

[0057] Another subject of the present invention is the use of the compounds of formula (I) for the preparation of H₃-antagonist medicaments according to the above-mentioned forms.

[0058] The invention further relates to the use of the compounds of formula (I) for preparing medicaments having the pre-cited effects.

30 [0059] Still another subject of the invention is a method for the treatment of precited ailments comprising administering a therapeutically effective dose of a compound (I), optionally in combination with a therapeutically acceptable vehicle or excipient.

[0060] For each of the above-indications, the amount of the active ingredient will depend upon the condition of the patient.

35 [0061] However, a suitable effective dose will be in general in the range of from 10 to 500 mg per day and of from 1 to 10 mg/day for particularly active compounds.

[0062] These doses are given on the basis of the compound and should be adapted for the salts, hydrates or hydrated salts thereof.

[0063] The invention is now illustrated by the following examples.

40

EXAMPLES






[0064] The structure of the synthesized compounds and their method of preparation as well as their melting point, recrystallisation solvent and elemental analysis are summarized in the following Table I:






45






50







55






TABLE I:






N°	FORMULA STRUCTURE NAME	mp (recryst. solv)	analysis (calc.)	method
1	$C_{16}H_{25}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-piperidine hydrogen oxalate	143-145°C (absolute ethanol)	C: 64.06 (64.07) H: 8.09 (8.16) N: 4.14 (4.15)	A
2	$C_{15}H_{23}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-pyrrolidine hydrogen oxalate	153-155°C (absolute ethanol)	C: 63.06 (63.14) H: 7.78 (7.79) N: 4.42 (4.33)	A
3	$C_{14}H_{23}NO; C_2H_2O_4$  N-methyl-N-(5-phenoxypropyl)-ethylamine hydrogen oxalate	122-124°C (absolute ethanol)	C: 61.74 (61.72) H: 8.24 (8.09) N: 4.52 (4.50)	A
4	$C_{15}H_{23}NO_2; C_2H_2O_4$  1-(5-phenoxypropyl)-morpholine hydrogen oxalate	166-168°C (absolute ethanol)	C: 60.10 (60.16) H: 7.45 (7.31) N: 4.08 (4.13)	A
5	$C_{17}H_{27}NO; C_2H_2O_4$  N-(5-phenoxypropyl)-hexamethyleneimine hydrogen oxalate	132-134°C (absolute ethanol)	C: 64.70 (64.93) H: 8.34 (8.32) N: 3.85 (3.99)	A







6	$C_{16}H_{27}NO; C_2H_2O_4$  N-ethyl-N-(5-phenoxypropyl)-propylamine hydrogen oxalate	90-91°C (isopropyl alcohol)	C: 63.60 (63.69) H: 8.81 (8.61) N: 3.97 (4.13)	B
7	$C_{17}H_{27}NO; 1.1 C_2H_2O_4$  1-(5-phenoxypropyl)-2-methyl-piperidine hydrogen oxalate	80-83°C (isopropyl alcohol)	C: 64.15 (63.98) H: 8.42 (8.17) N: 3.97 (3.89)	B
8	$C_{19}H_{31}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-4-propyl-piperidine hydrogen oxalate	165-166°C (absolute ethanol)	C: 66.27 (66.46) H: 8.94 (8.76) N: 3.72 (3.69)	B
9	$C_{17}H_{27}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-4-methyl-piperidine hydrogen oxalate	151-152°C (absolute ethanol)	C: 64.87 (64.93) H: 8.41 (8.32) N: 4.01 (3.99)	B
10	$C_{17}H_{27}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-3-methyl-piperidine hydrogen oxalate	140-141°C (isopropyl alcohol)	C: 65.35 (64.93) H: 8.49 (8.32) N: 4.00 (3.99)	B






11	$C_{17}H_{25}N_2O_2; C_2H_2O_4$  1-acetyl-4-(5-phenoxy)pentyl-piperazine hydrogen oxalate	186-188°C (absolute ethanol)	C: 59.78 (59.99) H: 7.47 (7.42) N: 7.35 (7.36)	B
12	$C_{18}H_{29}NO; 1.05 C_2H_2O_4$  1-(5-phenoxy)pentyl-3,5-trans-dimethyl- piperidine hydrogen oxalate	154-155°C (absolute ethanol)	C: 65.16 (65.25) H: 8.61 (8.47) N: 3.66 (3.79)	B
13	$C_{18}H_{29}NO; C_2H_2O_4$  1-(5-phenoxy)pentyl-3,5-cis-dimethyl- piperidine hydrogen oxalate	154-155°C (isopropyl alcohol)	C: 65.62 (65.73) H: 8.64 (8.55) N: 3.63 (3.83)	B
14	$C_{18}H_{29}NO; HCl$  1-(5-phenoxy)pentyl-2,6-cis-dimethyl- piperidine hydrochloride	135-136°C (acetone)	C: 69.18 (69.32) H: 9.79 (9.70) N: 4.28 (4.49)	B
15	$C_{19}H_{29}NO_3; C_2H_2O_4$  4-carboethoxy-1-(5-phenoxy)pentyl- piperidine hydrogen oxalate	149-150°C (absolute ethanol)	C: 61.16 (61.60) H: 7.76 (7.63) N: 3.40 (3.42)	B

16	$C_{19}H_{29}NO_3; C_2H_2O_4$  3-carboethoxy-1-(5-phenoxy)pentyl-piperidine hydrogen oxalate	117-118°C (isopropyl alcohol)	C: 61.54 (61.60) H: 7.87 (7.63) N: 3.29 (3.42)	B
17	$C_{16}H_{23}NO; C_2H_2O_4$  1-(5-phenoxy)pentyl-1,2,3,6-tetrahydropyridine hydrogen oxalate	177-179°C (methanol)	C: 64.19 (64.46) H: 7.49 (7.51) N: 4.25 (4.18)	B
18	$C_{15}H_{22}N_2O_3; C_2H_2O_4; 0.2 H_2O$  1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	145-147°C (absolute ethanol)	C: 54.89 (54.89) H: 6.68 (6.61) N: 7.41 (7.53)	C
19	$C_{15}H_{22}ClNO; C_2H_2O_4$  1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	139-141°C (absolute ethanol)	C: 57.00 (57.06) H: 6.63 (6.76) N: 3.79 (3.91) Cl: 10.24 (9.91)	C
20	$C_{16}H_{25}NO_2; C_2H_2O_4$  1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	115-116°C (absolute ethanol)	C: 61.22 (61.17) H: 7.72 (7.70) N: 4.03 (3.96)	C
21	$C_{16}H_{25}NO; C_2H_2O_4$  1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	138-140°C (absolute ethanol)	C: 64.05 (64.07) H: 8.00 (8.07) N: 4.10 (4.15)	C






22	$C_{16}H_{22}N_2O$; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	129-130°C (absolute ethanol)	C: 61.24 (61.16) H: 6.81 (6.82) N: 7.95 (7.84)	C
23	$C_{19}H_{25}NO$; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine hydrogen oxalate	166-167°C (methanol)	C: 67.42 (67.54) H: 7.26 (7.29) N: 3.66 (3.75)	C
24	$C_{19}H_{25}NO$; 1.25 $C_2H_2O_4$  1.25 $(COOH)_2$ 1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine hydrogen oxalate	160-163°C (methanol)	C: 65.12 (65.22) H: 7.17 (7.00) N: 3.52 (3.54)	C
25	$C_{15}H_{22}ClNO$; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	131-132°C (absolute ethanol)	C: 56.94 (57.06) H: 6.67 (6.76) N: 3.74 (3.91) Cl: 9.64 (9.91)	C
26	$C_{21}H_{27}NO$; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	189-190°C (methanol)	C: 69.16 (69.15) H: 7.39 (7.32) N: 3.39 (3.51)	C






27	$C_{19}H_{29}NO$; $C_2H_2O_4$  1-[5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine hydrogen oxalate	131-132°C (absolute ethanol)	C: 66.73 (66.82) H: 8.37 (8.28) N: 3.68 (3.71)	C
28	$C_{21}H_{27}NO$; 1.1 $C_2H_2O_4$  1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	155-157°C (absolute ethanol)	C: 68.40 (68.22) H: 7.04 (7.21) N: 3.45 (3.43)	C
29	$C_{15}H_{21}NO$; $C_2H_2O_4$  1-(5-phenoxy)pentyl-2,5-dihydropyrrole hydrogen oxalate	140-141°C (absolute ethanol)	C: 63.45 (63.54) H: 7.26 (7.21) N: 4.26 (4.36)	B
30	$C_{19}H_{29}NO$; $C_2H_2O_4$  1-[5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 66.99 (66.82) H: 8.47 (8.28) N: 3.72 (3.71)	C
31	$C_{14}H_{21}NO$; $C_2H_2O_4$  1-(4-phenoxybutyl)-pyrrolidine hydrogen oxalate	143-144°C (absolute ethanol)	C: 62.25 (62.12) H: 7.46 (7.49) N: 4.49 (4.53)	C







32	$C_{16}H_{25}NO$; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-(6-phenoxyhexyl)-pyrrolidine hydrogen oxalate	146-147°C (absolute ethanol)	C: 63.06 (63.10) H: 8.03 (7.91) N: 4.32 (4.04)	C
33	$C_{15}H_{23}NS$; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-(5-phenylthiopentyl)-pyrrolidine hydrogen oxalate	150-152°C (absolute ethanol)	C: 59.52 (59.29) H: 7.44 (7.29) N: 4.06 (4.02)	C
34	$C_{14}H_{21}NS$; $C_2H_2O_4$  $(COOH)_2$ 1-(4-phenylthiobutyl)-pyrrolidine hydrogen oxalate	114-116°C (absolute ethanol)	C: 59.24 (59.05) H: 7.16 (7.12) N: 4.16 (4.30) S: 9.79 (9.85)	C
35	$C_{13}H_{19}NO$; $C_2H_2O_4$  $(COOH)_2$ 1-(3-phenoxypropyl)-pyrrolidine hydrogen oxalate	169-170°C (absolute ethanol)	C: 60.98 (61.00) H: 7.14 (7.17) N: 4.64 (4.74)	C
36	$C_{15}H_{22}N_2O_3$; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	130-131°C (absolute ethanol)	C: 55.30 (55.43) H: 6.55 (6.57) N: 7.49 (7.60)	C
37	$C_{15}H_{22}FNO$; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	149-150°C (absolute ethanol)	C: 59.52 (59.81) H: 7.12 (7.09) N: 4.05 (4.10)	C






38	$C_{17}H_{26}N_2O_3; C_2H_2O_4$  1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 57.32 (57.55) H: 7.19 (7.12) N: 6.89 (7.07)	C
39	$C_{17}H_{25}NO_2; C_2H_2O_4$  1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	130-134°C (absolute ethanol)	C: 62.43 (62.45) H: 7.41 (7.45) N: 3.75 (3.83)	D
40	$C_{15}H_{24}N_2O; 2.1 C_2H_2O_4$  1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine di-(hydrogen oxalate)	120-122°C (absolute ethanol)	C: 52.49 (52.72) H: 6.74 (6.50) N: 6.32 (6.40)	E1
41	$C_{16}H_{22}N_2O; C_2H_2O_4$  1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	119-120°C (absolute ethanol)	C: 61.95 (62.05) H: 6.88 (6.94) N: 8.00 (8.04)	C
42	$C_{13}H_{20}N_2O_3; C_2H_2O_4$  N-[3-(4-nitrophenoxy)-propyl]-diethylamine hydrogen oxalate	160-161°C (absolute ethanol/ methanol 1:1)	C: 52.46 (52.63) H: 6.49 (6.48) N: 8.10 (8.12)	F

43	$C_{14}H_{20}N_2O_2$; $C_2H_2O_4$ N-[3-(4-cyanophenoxy)-propyl]-diethylamine hydrogen oxalate	148-150°C (absolute ethanol)	C: 59.40 (59.62) H: 6.82 (6.88) N: 8.60 (8.69)	F
44	$C_{22}H_{27}NO_2$; $C_2H_2O_4$ 1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	141-142°C (absolute ethanol)	C: 67.17 (67.43) H: 6.80 (6.84) N: 3.18 (3.28)	D
45	$C_{23}H_{29}NO_2$; $C_2H_2O_4$ 1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine hydrogen oxalate	177-178°C (absolute ethanol)	C: 67.77 (68.01) H: 7.09 (7.08) N: 3.26 (3.17)	D
46	$C_{15}H_{23}NO_2$; 1.1 $C_2H_2O_4$ N-[3-(4-acetylphenoxy)-propyl]-diethylamine hydrogen oxalate	108-110°C (absolute ethanol)	C: 59.30 (59.30) H: 7.47 (7.29) N: 4.18 (4.02)	F
47	$C_{17}H_{26}N_2O_2$; $C_2H_2O_4$ 1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	142-144°C (absolute ethanol)	C: 59.67 (59.99) H: 7.55 (7.42) N: 7.25 (7.36)	C



48	$C_{21}H_{27}NO_2$; $C_2H_2O_4$  1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	135-136°C (absolute ethanol)	C: 66.49 (66.49) H: 7.05 (7.04) N: 3.24 (3.37)	D
49	$C_{22}H_{28}N_2O_2$; 1.1 $C_2H_2O_4$  1.1 ($COOH$) ₂ 1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	176-178°C (absolute ethanol)	C: 64.56 (64.38) H: 6.89 (6.74) N: 6.26 (6.20)	E ₂
50	$C_{17}H_{27}NO_2$; $C_2H_2O_4$  1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]- pyrrolidine hydrogen oxalate	102-104°C (absolute ethanol)	C: 61.89 (62.11) H: 7.94 (7.96) N: 3.77 (3.81)	G
51	$C_{16}H_{24}N_2O$; $C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-diethylamine hydrogen oxalate	120-122°C (absolute ethanol)	C: 61.56 (61.70) H: 7.54 (7.48) N: 7.87 (7.99)	H
52	$C_{17}H_{24}N_2O$; $C_2H_2O_4$  1-[5-(4-cyanophenoxy)-pentyl]-piperidine hydrogen oxalate	115-116°C (absolute ethanol)	C: 62.62 (62.97) H: 7.20 (7.23) N: 7.76 (7.73)	H

53	$C_{14}H_{20}N_2O; C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine hydrogen oxalate	148-149°C (absolute ethanol)	C: 59.68 (59.62) H: 6.76 (6.88) N: 8.57 (8.69)	H
54	$C_{13}H_{18}N_2O; C_2H_2O_4$  N-[2-(4-cyanophenoxy)-ethyl]-diethylamine hydrogen oxalate	124-125°C (absolute ethanol)	C: 58.15 (58.43) H: 6.30 (6.54) N: 8.95 (9.09)	H
55	$C_{12}H_{16}N_2O; C_2H_2O_4$  N-[3-(4-cyanophenoxy)-propyl]-dimethylamine hydrogen oxalate	166-167°C (absolute ethanol/ methanol 1:1)	C: 57.01 (57.14) H: 6.02 (6.16) N: 9.46 (9.52)	H
56	$C_{15}H_{22}N_2O; C_2H_2O_4$  N-[4-(4-cyanophenoxy)-butyl]-diethylamine hydrogen oxalate	143-145°C (absolute ethanol)	C: 60.80 (60.70) H: 7.11 (7.19) N: 8.22 (8.33)	H
57	$C_{18}H_{28}N_2O; C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine hydrogen oxalate	134-136°C (absolute ethanol)	C: 63.38 (63.47) H: 8.11 (7.99) N: 7.29 (7.40)	H

58	$C_{14}H_{18}N_2O$; 1.1 $C_2H_2O_4$  1.1 (COOH) ₂ 1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine hydrogen oxalate	163-165°C (absolute ethanol)	C: 58.95 (59.08) H: 6.23 (6.18) N: 8.43 (8.51)	H
59	$C_{15}H_{20}N_2O$; 1.05 $C_2H_2O_4$  1.05 (COOH) ₂ 1-[3-(4-cyanophenoxy)-propyl]-piperidine hydrogen oxalate	151-153°C (absolute ethanol)	C: 60.62 (60.61) H: 6.66 (6.57) N: 8.25 (8.27)	H
60	$C_{16}H_{22}N_2O$; 1.05 $C_2H_2O_4$  1.05 (COOH) ₂ N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine hydrogen oxalate	124-125°C (absolute ethanol)	C: 61.62 (61.60) H: 6.94 (6.88) N: 7.87 (7.94)	H
61	$C_{17}H_{26}N_2O$; $C_2H_2O_4$  (COOH) ₂ N-[6-(4-cyanophenoxy)-hexyl]-diethylamine hydrogen oxalate	110-112°C (absolute ethanol)	C: 62.90 (62.62) H: 7.76 (7.74) N: 7.61 (7.69)	H
62	$C_{16}H_{24}N_2O$; $C_2H_2O_4$  (COOH) ₂ N-[3-(4-cyanophenoxy)-propyl]-dipropylamine hydrogen oxalate	127-128°C (absolute ethanol)	C: 61.57 (61.70) H: 7.57 (7.48) N: 7.91 (7.99)	H
63	$C_{15}H_{25}NO_2$; $C_2H_2O_4$; 0.5 H_2O  (COOH) ₂ 0.5 H_2O N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine hydrogen oxalate hemihydrate	33-36°C (isopropyl alcohol)	C: 58.15 (58.27) H: 8.15 (8.05) N: 4.21 (4.00)	G

64	$C_{15}H_{24}N_2O_2$; $C_2H_2O_4$  4'-(3-diethylaminopropoxy)-acetophenone-oxime hydrogen oxalate	99-100°C (absolute ethanol)	C: 57.26 (57.61) H: 7.47 (7.39) N: 7.72 (7.90)	J
65	$C_{16}H_{23}NO_2$; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-piperidine hydrogen oxalate	159-160°C (absolute ethanol)	C: 61.18 (61.52) H: 7.11 (7.17) N: 3.96 (3.99)	K
66	$C_{17}H_{25}NO_2$; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine hydrogen oxalate	143-144°C (absolute ethanol)	C: 62.11 (62.45) H: 7.41 (7.45) N: 3.79 (3.83)	K
67	$C_{18}H_{27}NO_2$; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl- piperidine hydrogen oxalate	171-172°C (absolute ethanol)	C: 63.06 (63.31) H: 7.44 (7.70) N: 3.64 (3.69)	K
68	$C_{17}H_{25}NO_2$; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine hydrogen oxalate	160-161°C (absolute ethanol)	C: 62.47 (62.45) H: 7.46 (7.45) N: 3.77 (3.83)	K

69	$C_{17}H_{25}NO_2$; $C_2H_2O_4$ 1-[3-(4-propionylphenoxy)-propyl]-piperidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 62.54 (62.45) H: 7.51 (7.45) N: 3.79 (3.83)	L
70	$C_{18}H_{27}NO_2$; $C_2H_2O_4$ 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl- piperidine hydrogen oxalate	174-175°C (absolute ethanol)	C: 63.22 (63.31) H: 7.60 (7.70) N: 3.64 (3.69)	K
71	$C_{15}H_{21}NO_2$; $C_2H_2O_4$ 1-[3-(4-formylphenoxy)-propyl]-piperidine hydrogen oxalate	152-153°C (absolute ethanol)	C: 60.23 (60.52) H: 6.81 (6.87) N: 4.15 (4.15)	L
72	$C_{18}H_{27}NO_2$; $C_2H_2O_4$ 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine hydrogen oxalate	121-122°C (absolute ethanol)	C: 63.02 (63.31) H: 7.73 (7.70) N: 3.66 (3.69)	L
73	$C_{16}H_{25}NO_2$; 1.5 $C_2H_2O_4$ N-[3-(4-propionylphenoxy)-propyl]-diethylamine hydrogen oxalate	118-120°C (absolute ethanol)	C: 57.27 (57.28) H: 7.00 (7.08) N: 3.47 (3.52)	L

74	$C_{18}H_{27}NO_2; C_2H_2O_4$  1-[3-(4-butyrylphenoxy)-propyl]-piperidine hydrogen oxalate	138-139°C (absolute ethanol)	C: 63.09 (63.31) H: 7.78 (7.70) N: 3.75 (3.69)	L
75	$C_{16}H_{21}NO_2; 1.1 C_2H_2O_4$  1.1 (COOH) ₂ 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6- tetrahydropyridine hydrogen oxalate	143-144°C (absolute ethanol)	C: 61.21 (61.00) H: 6.25 (6.52) N: 4.00 (3.91)	K

[0065] Compounds 1 to 75 are prepared according to the following procedures:

METHOD A:

[0066] A solution of 1-bromo-5-phenoxy-pentane (1.4 to 3.5 mmol) in ten equivalents of the suitable secondary amine was heated to reflux temperature with stirring for 48 hours (compds. 1, 3 and 4), 24 hours (compd. 2) or 4 hours (compd. 5). After cooling, the excess base was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate formed was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD B:

[0067] A solution of 1-bromo-5-phenoxy-pentane (0.9 to 1.7 mmol) and an excess of the suitable secondary amine (2.3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 48 hours (compd. 6) or 24 hours (compds. 7, 8, 9, 10, 11, 12&13, 14, 15, 16, 17 and 29). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 12 and 13 were separated by column chromatography on silica gel eluting with a solvent mixture of petroleum spirit (bp 60-80°C), diethyl ether and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compounds 14 and 16 were purified by column chromatography on silica gel eluting with diethyl ether and triethylamine in the ratio 99:1, and the eluent was removed under reduced pressure to leave an oil. The oil was converted to oxalate salt (compds. 6, 7, 8, 9, 11, 12, 13, 15, 16, 17 and 29) by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from isopropyl alcohol (compds. 6, 7, 10, 13 and 16), absolute ethanol (compds. 8, 9, 11, 12, 15 and 29) or methanol (compd. 17). The oil was converted to hydrochloride salt (compd. 14) by adding 2N HCl. The precipitate was formed in a mixture of chloroform and diethyl ether (1:1) and recrystallised from acetone.

METHOD C:

[0068] A solution of the suitable α -bromo- ω -aryloxy alkane (0.4 to 1.4 mmol) or ω -bromoalkyl phenyl sulphide (1 mmol, compds. 33 and 34) and an excess of pyrrolidine (10 to 15 equivalents) or 3-methylpiperidine (10 equivalents,

compd. 38) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours or 16 hours (compd. 47). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD D:

[0069] A solution of the suitable 4'-(5-bromopentoxy)phenyl ketone (0.7 to 1 mmol, compds. 39, 44 and 45) or 1-bromo, 5-(4-phenoxyphenoxy)pentane (0.6 mmol, compd. 48) and an excess of pyrrolidine (10 to 15 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours (compds. 39, 44 and 48) or 24 hours (compd. 45). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with chloroform (compds. 39, 45 and 48) or dichloromethane (compd. 44), the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (recrystallised twice from absolute ethanol in the case of compd. 39).

METHOD E:

[0070]

1. The oxalate 18 was prepared according to method C. A solution of compound 18 (0.57 mmol) in 10 ml methanol and 10 ml absolute ethanol was placed with 100 mg of palladium (5%) on carbon catalyst in a two-neck round-bottom flask fitted with a balloon filled with hydrogen. The mixture was stirred vigorously at room temperature and the flask was purged of air and filled with hydrogen. After 3 hours, the catalyst was filtered off on celite and the solvent removed under reduced pressure. The residual solid was converted to oxalate salt by dissolving in methanol and adding a solution of oxalic acid (2 equivalents) in absolute ethanol. Diethyl ether was added to form a precipitate. The product was recrystallised from absolute ethanol.

2. To a solution of compound 40 (0.35 mmol) in pyridine vigorously stirred at 0°C was added dropwise a slight excess of benzoyl chloride (0.4 mmol). The stirring was allowed to continue 20 minutes after the end of the addition after which the mixture was placed in the refrigerator overnight (16 hours). The solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with chloroform, the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was dissolved in methanol, filtered, and concentrated under reduced pressure the solid was recrystallised from absolute ethanol.

METHOD F:

[0071] In a three-neck flask kept under nitrogen was placed a solution of the suitable phenol (1.6 mmol), 3-(diethyl-amino)propanol (1.5 mmol), and triphenyl phosphine (1.9 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diisopropyl azodicarboxylate (2 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCl (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (compds. 43 and 46) or from a 1:1 mixture of methanol and absolute ethanol (compd. 42).

METHOD G:

[0072] A solution of the free base of compound 39 (0.6 mmol) or compound 46 (0.8 mmol) in 20 ml dry diethyl ether

was added dropwise to a stirred suspension of lithium aluminium hydride (0.6 or 0.8 mmol) in 20 ml dry diethyl ether kept under nitrogen. The mixture was stirred at room temperature under nitrogen for two hours. Ice-cold water was carefully added and the organic layer decanted. The aqueous phase was extracted with diethyl ether. The combined organic solutions were dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a yellow oil.

5 The oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (compd 50) or from isopropyl alcohol, giving a very hygroscopic solid (compd. 63).

METHOD H:

10 **[0073]** A solution of the suitable α -bromo- ω -(4-cyanophenoxy) alkane (0.5 to 0.7 mmol) and an excess of the suitable secondary amine (8 to 12 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours (compds. 54, 55, 57 and 60), 20 hours (compd. 52), 16 hours (compds. 56, 58, 59 and 61) or 8 hours (compd. 51) or was stirred at room temperature for 48 hours (compd. 53) or 24 hours (compd. 60). After cooling, the solvent was

15 removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 62 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. For all the compounds of method H, the remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in

20 absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (two recrystallisations were required for compds. 58 and 59) or from a 1:1 mixture of methanol and absolute ethanol (compd. 55).

METHOD J:

25 **[0074]** A solution of compound 46 (1 mmol) in 10 ml methanol was stirred at room temperature and a solution of hydroxylamine hydrochloride (2 equivalents) in 2 ml water was added. The mixture was stirred at 50-70°C in a water bath for 20 minutes. Methanol was removed under reduced pressure. The residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 64 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. Diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD K:

35 **[0075]** A solution of 4'-(3-bromopropoxy)acetophenone (0.8 to 1.9 mmol) and an excess of the suitable piperidine (3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours. After cooling,

40 the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 67 and 70 were separated by column chromatography on silica gel eluting with a solvent mixture of diethyl ether, petroleum spirits (bp 60-80°C) and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compound 75 was purified by column chromatography on silica gel eluting with chloroform and methanol (1:1), and concentrated under reduced pressure.

45 The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

METHOD L:

50 **[0076]** In a three-neck flask kept under nitrogen was placed a solution of the suitable 4'-hydroxyphenyl ketone (0.9 to 3 mmol), 3-(1-piperidinyl)propanol (0.9 to 3 mmol), and triphenyl phosphine (1 to 3.5 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diethyl azodicarboxylate (1 to 3.6 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCl (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane.

ane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The crude product was purified by column chromatography on silica gel eluting with diethyl ether containing 1 % triethylamine, and concentrated under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

Pharmacological study

[0077] Interaction of compounds with the H_3 receptor are evidenced *in vitro* by the measurement of the release of neosynthesized tritiated histamine from rat cerebral cortex synaptosomes preincubated with tritiated histidine (Garbarg et al., J. Pharmacol. Exp. Ther., 1992, 263: 304-310). The H_3 potency of compounds is measured by the progressive reversal of the tritiated histamine release inhibition by the selective H_3 agonist (R) α -methylhistamine (Arrang et al., Nature, 1987, 327: 117-123).


[0078] The effects of antagonists were estimated *in vivo* by the measurement of the tele-methylhistamine level variations in the brain of mice (Garbarg et al., J. Neurochem., 1989, 53: 1724-1730). At various time after p.o. administration of the compound, the effect of the H_3 antagonist is evidenced by the increase in the telemethylhistamine level induced. This increase is compared to the maximal effect induced by the reference H_3 -antagonist thioperamide given at the dose of 10 mg/kg, p.o. This allows the calculation of the ED_{50} value for each compound which correspond to the dose responsible for an half maximal effect.

[0079] The results are reported in the following table II:

Ex No.	X	n	R ¹ R ²	R ³ (n ₃ = 1)	Ki(nM)	ED ₅₀ (mg/kg/p.o.)
18	O	5	-(CH ₂) ₄ -	p-NO ₂	39±11	1.1
43	O	3	Et, Et	p-CN	95±28	0.50
46	O	3	Et, Et	p-CH ₃ CO		0.44
50	O	5	-(CH ₂) ₄ -	p-CH ₃ CH(OH)		1.0
56	O	4	Et, Et	p-CN		1.1
59	O	3	-(CH ₂) ₅ -	p-CN		0.20
60	O	3	-(CH ₂) ₆ -	p-CN		0.64
63	O	3	Et, Et	p-CH ₃ CH(OH)		0.34
64	O	3	Et, Et	p-CH ₃ C=N(OH)		0.8
66	O	3	-(3-Me)-(CH ₂) ₅ -	p-CH ₃ CO		0.3
68	O	3	-(4-Me)-(CH ₂) ₅ -	p-CH ₃ CO		0.3
69	O	3	-(CH ₂) ₅ -	p-C ₂ H ₅ CO		0.4

Claims

1. Compound of general formula (I) in which:




- C_nH_{2n} is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- R^1 and R^2 may be identical or different and represent each independently

- a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
- a saturated nitrogen-containing ring



- with m ranging from 4 to 7, or
- an unsaturated nitrogen-containing ring



- with p, q and r being 1 to 3 independently, such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or
- a morpholino group, or
- a N-substituted piperazino group:



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- n_3 is an integer from 0 to 5;
- R^3 represents each independently

- a halogen atom,
- a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkylloximino, aryloximino, α -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
- or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring,

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, except compounds in which

- $-NR^1R^2$ is a pyrrolidinyl group, C_nH_{2n} is a linear chain $-(CH_2)_n-$ and n_3 is zero, X being an oxygen atom with n ranging from 3 to 5, or X being a sulfur atom with n being 4 or 5;
- $-NR^1R^2$ is a piperidinyl group, C_nH_{2n} is a linear chain $-(CH_2)_n-$ and X is an oxygen atom, n_3 being zero with n being 2, 5 or 8 or n_3 being 1 with R^3 being 4-CN and n being 5;
- $-NR^1R^2$ is a diethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain $-(CH_2)_n-$ and n_3 is 1, R^3 being 4-

NO₂ or 4-COCH₃ with n being 3 or R³ being 4-CN with n being 2 to 4;


- -NR¹R² is a dimethylamine group, X is an oxygen atom, C_nH_{2n} is a linear chain -(CH₂)_n- and n³ is 1, R³ being 4-CN with n being 3.

- 5 2. Compound according to claim 1, in which R¹ and R² are independently a lower alkyl group.
3. Compound according to claim 2, in which R¹ and R² are each an ethyl group.
- 10 4. Compound according to claim 1, in which -NR¹R² is a saturated nitrogen-containing ring: m being as



defined in claim 1.

- 20 5. Compound according to claim 4, characterized in that m is 4, 5 or 6.
6. Compound according to claim 1, characterized in that -NR¹R² is an unsaturated nitrogen-containing ring:



p, q and r being as defined in claim 1, preferably p, q and r are 1 or 2, more preferably p is 2 and q and r are 1.

- 35 7. Compound according to anyone of claims 4 to 6, characterized in that the nitrogen-containing ring i) or ii) is unsubstituted.
8. Compound according to anyone of claim 4 to 6, characterized in that the nitrogen-containing ring i) or ii) is substituted, preferably mono-substituted with an alkyl group.
- 40 9. Compound according to claim 8, characterized in that the nitrogen-containing ring is mono-substituted with a methyl group.
10. Compound according to claim 1, characterized in that -NR¹R² is a morpholino group.
- 45 11. Compound according to claim 1, characterized in that -NR¹R² is a N-substituted piperazino group, preferably N-acetylpiperazino.
12. Compound according to anyone of claims 1 to 11, characterized in that n₃ is zero.
- 50 13. Compound according to anyone of claims 1 to 11, characterized in that n₃ is 1 with R³ being as defined in claim 1 and preferably in para-position.
14. Compound according to anyone of claims 1 to 11 and 13, characterized in that R³ is a lower alkyl, preferably a C₁-C₄ alkyl.
- 55 15. Compound according to anyone of claims 1 to 11 and 13, characterized in that R³ is a halogen atom, a cyano, nitro, alkanoyl, alkyloximine or hydroxyalkyl, preferably CN, NO₂, COCH₃, COC₂H₅, H₃C-C=N-OH or H₃C-CHOH.

16. Compound according to anyone of claims 1 to 11, characterized in that R³ taken together with the carbon atoms of the phenyl group to which it is fused, form a 5- or 6- membered saturated or unsaturated ring, in particular a 5,6,7,8-tetrahydronaphthyl group.

17. Compound according to anyone of claims 1 to 11, characterized in that R³ taken together with the phenyl group to which it is fused, form a naphthyl group.

18. Compound according to anyone of claims 1 to 17, characterized in that -C_nH_{2n}- is a linear hydrocarbon chain - (CH₂)_n-, n being as defined in claim 1.

19. Compound according to anyone of claims 1 to 18, characterized in that X is an oxygen atom.

20. Compound according to anyone of claims 1 to 18, characterized in that X is a sulfur atom.

21. Compound according to anyone of claims 1 to 20, characterized in that n is varying from 3 to 5 and is preferably 3.

22. Compound according to anyone of claims 1 to 21, characterized in that it is one of the following compounds:

N-methyl-N-(5-phenoxy-pentyl)-ethylamine
 1-(5-phenoxy-pentyl)-morpholine
 N-(5-phenoxy-pentyl)-hexamethyleimine
 N-ethyl-N-(5-phenoxy-pentyl)-propylamine
 1-(5-phenoxy-pentyl)-2-methyl-piperidine
 1-(5-phenoxy-pentyl)-4-propyl-piperidine
 1-(5-phenoxy-pentyl)-4-methyl-piperidine
 1-(5-phenoxy-pentyl)-3-methyl-piperidine
 1-acetyl-4-(5-phenoxy-pentyl)-piperazine
 1-(5-phenoxy-pentyl)-3,5-trans-dimethyl-piperidine
 1-(5-phenoxy-pentyl)-3,5-cis-dimethyl-piperidine
 1-(5-phenoxy-pentyl)-2,6-cis-dimethyl-piperidine
 4-carboethoxy-1-(5-phenoxy-pentyl)-piperidine
 3-carboethoxy-1-(5-phenoxy-pentyl)-piperidine
 1-(5-phenoxy-pentyl)-1,2,3,6-tetrahydropyridine
 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
 1-[5-(2-naphthyl-oxy)-pentyl]-pyrrolidine
 1-[5-(1-naphthyl-oxy)-pentyl]-pyrrolidine
 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
 1-[5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
 1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
 1-(5-phenoxy-pentyl)-2,5-dihydropyrrole
 1-[5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
 1-(6-phenoxyhexyl)-pyrrolidine
 1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
 1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
 1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
 1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine
 1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine

1-[5-(4-(1-hydroxyethyl)-phenoxy)-pentyl]-pyrrolidine
 1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
 N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
 N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
 1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
 N-[6-(4-cyanophenoxy)-hexyl]-diethylamine
 N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
 1-[3-(4-formylphenoxy)-propyl]-piperidine
 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
 N-[3-(4-propionylphenoxy)-propyl]-diethylamine
 1-[3-(4-butyrylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine


23. Compound according to anyone of claims 1 to 22, characterized in that it is one of the following compounds:

1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine

24. Pharmaceutical composition characterized in that it comprises as active ingredient, a therapeutically effective amount of a compound according to anyone of claim 1 to 23 in combination with a pharmaceutically acceptable vehicle or excipient.


25. Medicament acting as an antagonist of the histamine H₃-receptors, characterized in that it comprises as active ingredient, an effective amount of a compound according to anyone of claims 1 to 23.

26. Use of a compound of general formula (I) in which:




- C_nH_{2n} is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- R¹ and R² may be identical or different and represent each independently

- a lower alkyl or cycloalkyl, or taken together with the nitrogen atom to which they are attached,
- a saturated nitrogen-containing ring



- with m ranging from 4 to 7, or
- an unsaturated nitrogen-containing ring



- with p, q and r being 1 to 3 independently, such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or
- a morpholino group, or
 - a N-substituted piperazino group:



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- n_3 is an integer from 0 to 5;
- R^3 represents each independently

- a halogen atom,
- a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkylloximino, aryloximino, α -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
- or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring.

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, for the preparation of a medicament acting as an antagonist of the histamine H_3 -receptors.

27. Use according to claim 26, characterized in that compound (I) is as defined in any one of claims 2 to 21.

28. Use according to claim 26 characterized in that compound (I) is one of the following compounds:

- 1-(5-phenoxypropyl)-piperidine
- 1-(5-phenoxypropyl)-pyrrolidine
- N-methyl-N-(5-phenoxypropyl)-ethylamine
- 1-(5-phenoxypropyl)-morpholine
- N-(5-phenoxypropyl)-hexamethyleneimine
- N-ethyl-N-(5-phenoxypropyl)-propylamine
- 1-(5-phenoxypropyl)-2-methyl-piperidine

1-(5-phenoxypropyl)-4-propyl-piperidine
 1-(5-phenoxypropyl)-4-methyl-piperidine
 1-(5-phenoxypropyl)-3-methyl-piperidine
 1-acetyl-4-(5-phenoxypropyl)-piperazine
 5 1-(5-phenoxypropyl)-3,5-trans-dimethyl-piperidine
 1-(5-phenoxypropyl)-3,5-cis-dimethyl-piperidine
 1-(5-phenoxypropyl)-2,6-cis-dimethyl-piperidine
 4-carboethoxy-1-(5-phenoxypropyl)-piperidine
 3-carboethoxy-1-(5-phenoxypropyl)-piperidine
 10 1-(5-phenoxypropyl)-1,2,3,6-tetrahydropyridine
 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
 15 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
 1-[5-(2-naphthyl)-pentyl]-pyrrolidine
 1-[5-(1-naphthyl)-pentyl]-pyrrolidine
 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
 20 1-[5-(2-(5,6,7,8-tetrahydronaphthyl)-oxy)-pentyl]-pyrrolidine
 1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
 1-(5-phenoxypropyl)-2,5-dihydropyrrole
 1-[5-(1-(5,6,7,8-tetrahydronaphthyl)-oxy)-pentyl]-pyrrolidine
 1-(4-phenoxybutyl)-pyrrolidine
 25 1-(6-phenoxyhexyl)-pyrrolidine
 1-(5-phenylthiopentyl)-pyrrolidine
 1-(4-phenylthiobutyl)-pyrrolidine
 1-(3-phenoxypropyl)-pyrrolidine
 1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
 30 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
 1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
 1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
 35 N-[3-(4-nitrophenoxy)-propyl]-diethylamine
 N-[3-(4-cyanophenoxy)-propyl]-diethylamine
 1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-(phenylacetyl)-phenoxy)-pentyl]-pyrrolidine
 N-[3-(4-acetylphenoxy)-propyl]-diethylamine
 40 1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
 1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
 1-[5-(4-(1-hydroxyethyl)-phenoxy)-pentyl]-pyrrolidine
 1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
 45 1-[5-(4-cyanophenoxy)-pentyl]-piperidine
 N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
 N-[2-(4-cyanophenoxy)-ethyl]-diethylamine
 N-[3-(4-cyanophenoxy)-propyl]-dimethylamine
 N-[4-(4-cyanophenoxy)-butyl]-diethylamine
 50 N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
 1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethylenimine
 N-[6-(4-cyanophenoxy)-hexyl]-diethylamine
 55 N-[3-(4-cyanophenoxy)-propyl]-dipropylamine
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-piperidine

1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine
 1-[3-(4-formylphenoxy)-propyl]-piperidine
 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine
 N-[3-(4-propionylphenoxy)-propyl]-diethylamine
 1-[3-(4-butyrylphenoxy)-propyl]-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine

29. Use according to claim 26, characterized in that compound (I) is one of the following compounds:

1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
 N-[3-(4-cyanophenoxy)-propyl]-diethylamine
 N-[3-(4-acetylphenoxy)-propyl]-diethylamine
 1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine
 N-[4-(4-cyanophenoxy)-butyl]-diethylamine
 1-[3-(4-cyanophenoxy)-propyl]-piperidine
 N-[3-(4-cyanophenoxy)-propyl]-hexamethylenimine
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine
 4-(3-diethylaminopropoxy)-acetophenone-oxime
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine
 1-[3-(4-propionylphenoxy)-propyl]-piperidine

30. Medicament according to anyone of claims 25 to 29, for the treatment of central nervous system disorders, in particular Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.

31. Medicament according to anyone of claims 25 to 29, having psychotropic effects, promoting wakefulness, attention, memory and improving mood, intended to be used in particular in the treatment of Alzheimer disease and other cognitive disorders in aged persons, depressive or asthenic states.

32. Medicament according to anyone of claims 25 to 29, having nootropic effects, intended to be used in particular in treatment to stimulate attention and memorization capacity.

33. Medicament according to anyone of claims 25 to 29, for the treatment of obesity, vertigo and motion sickness.

34. Medicament according to anyone of claims 25 to 29, for the treatment of CNS disorders, in particular of aged persons.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 98 40 1944 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.8)
X	GB 1 512 880 A (MITSUBISHI CHEM IND) 1 June 1978 * examples; table I *	1-24	C07D295/088 C07C211/08 C07D211/04 C07D295/185
X	DE 12 69 134 B (VEB DEUTSCHES HYDRIERWERK RODLEREN) 18 December 1962 * examples 1-5 *	1-24	C07D211/62 C07D211/70 C07D207/20
X	US 3 947 434 A (SPENCER CLAUDE F ET AL) 30 March 1976 * examples LXXIXA, B *	1-24	A61K31/13 A61K31/40 A61K31/445 A61K31/495
X	US 4 751 302 A (IBUKI TADAYUKI ET AL) 14 June 1988 * tables 5-2, 5-3 *	1-24	
X	GB 924 961 A (THE WELLCOME FOUNDATION LIMITED) 1 May 1963 * table II *	1-24	
X	US 3 312 696 A (TURBANTI L.) 4 April 1967 * claim 1; examples 1-20 *	1-24	
	-/--		TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07D C07C A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search	Date of completion of the search	Examiner	
MUNICH	7 December 1998	Juntunen, A	
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 98 40 1944

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	DE 26 24 261 A (BOTTU FA) 16 December 1976 * page 4; example 10 * ---	1-24	
X	US 2 870 151 A (WRIGHT H.B. AND MOORE M.B.) 20 January 1959 * examples I-XII * ---	1-24	
X	DE 965 813 C (ABBOTT LABORATORIES) 19 June 1957 * examples 1-5,7 * ---	1-24	
X	LITTMANN E. R. AND MARVEL C. S.: "Cyclic Quaternary Ammonium Salts from Halogenated Aliphatic Tertiary Amines" J.AMER.CHEM.SOC., vol. 52, 1930, pages 287-294, XP002084866 * page 289 - page 290 * ---	1-24	
X	KIKUMOTO R. ET AL.: "Synthesis and Antidepressant Activity of Substituted (gamma-Aminoalkoxy)benzene Derivatives" J.MED.CHEM., vol. 24, no. 2, 1981, pages 145-148, XP000565653 * table 1 * ---	1-26	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	SHADBOLT R. S. ET AL.: "Some Aryloxyalkylamines, N-Arylethylenediamines and Related Compounds as Anorectic Agents" J.MED.CHEM., vol. 14, no. 9, 1971, pages 836-842, XP002084867 * table 1 * ---	1-24	
	---	-/--	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 98 40 1944

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	<p>WALSH D. A. ET AL.: "Synthesis and Antiallergy Activity of 4-(Diarylhydroxymethyl)-1-[3-(aryloxy)propyl]piperidines and Structurally Related Compounds"</p> <p>J.MED.CHEM., vol. 32, no. 1, 1989, pages 105-118, XP002084868 * tables I-III *</p> <p>---</p>	1-24	
X	<p>SOHDA T ET AL: "STUDIES ON ANTIDIABETIC AGENTS. SYNTHESIS OF 5-4-(1-METHYLCYCLOHEXYLMETHOXY)-BENZYL)THIAZOLIDINE-2,4-DIONE (ADD-3878) AND ITS DERIVATIVES"</p> <p>CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 30, no. 10, 1982, pages 3580-3600, XP002046259 * table VIII *</p> <p>---</p>	1-24	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X	<p>CHABRIER P. ET AL.: "Nouveaux carbamates doués d'activité anesthésique locale"</p> <p>BULL.SOC.CHIM.FR.,1955, pages 1353-1357, XP002084869 * table IV *</p> <p>---</p>	1-24	
X	<p>MARQUET J. ET AL.: "Topologically Controlled Coulombic Interactions, a New Tool in the Developing of Novel Reactivity. Photochemical and Electrochemical Cleavage of Phenyl Alkyl Ethers"</p> <p>J.ORG.CHEM., vol. 60, no. 12, 1995, pages 3814-3825, XP002084870 * table 1 *</p> <p>---</p>	1-24	
	---	-/--	



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 98 40 1944

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	CASANOVAS A.-M. ET AL.: "Etude des relations structure-activité d'une série d'anesthésiques locaux" EUR.J.MED.CHEM.-CHIM.THER., vol. 17, no. 4, 1982, pages 333-337, XP002084871 * page 334 *	1-24	
X	--- KIKUMOTO R ET AL: "SYNTHESES AND PLATELET AGGREGATION INHIBITORY AND ANTITHROMBOTIC PROPERTIES OF 2-(OMEGA-AMINOALKOXY)PHENYLETHYLBENZENES" JOURNAL OF MEDICINAL CHEMISTRY, vol. 33, no. 6, June 1990, pages 1818-1823, XP000673455 * tables I-III *	1-24	
X	--- CHENEY L.C. ET AL.: "Alkylaminoalkyl Ethers of the Benzylphenols" J.AMER.CHEM.SOC., vol. 71, 1949, pages 60-64, XP002086293 * page 60; table I *	1-26	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
A	--- STARK H. ET AL.: "Developments of Histamine H3-receptor Antagonists" DRUGS OF THE FUTURE, vol. 21, no. 5, 1996, pages 507-520, XP002084872 * page 507 *	1-34	



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 98 40 1944

Claim(s) searched incompletely:
1-34

Reason for the limitation of the search:

The search on the final compounds of a restricted subset of formula I (R1 and R2= a lower alkyl, a saturated N-containing ring, a morpholino group, a N-substituted piperazino group as defined in claim 1) and their histamine H3-receptor antagonistic activity revealed already a vast amount of novelty destroying compounds with respect to claim 1 of the present application. Therefore the search had to be limited to the compounds of claims 2 and 5 encompassed by the above defined subset, and to the activity thereof.

Despite the above limitation to the two groups of compounds the search revealed too many relevant documents and/or compounds so that the search report shall not be considered complete.

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 40 1944

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-12-1998

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1512880 A	01-06-1978	JP	1233459 C	26-09-1984
		JP	52000248 A	05-01-1977
		JP	59008265 B	23-02-1984
		US	4024282 A	17-05-1977
		CH	623301 A	29-05-1981
		DE	2627227 A	30-12-1976
		DK	276276 A,B,	20-12-1976
		FR	2315913 A	28-01-1977
		NL	7606668 A,B,	21-12-1976
		SE	430156 B	24-10-1983
		SE	7607013 A	20-12-1976
		US	4071559 A	31-01-1978
		JP	1258356 C	29-03-1985
		JP	52033635 A	14-03-1977
		JP	59035386 B	28-08-1984
		US	4061776 A	06-12-1977
		JP	1283612 C	27-09-1985
		JP	52033658 A	14-03-1977
		JP	60006349 B	18-02-1985
		US	4091114 A	23-05-1978
		JP	1323708 C	27-06-1986
		JP	52057133 A	11-05-1977
		JP	60048507 B	28-10-1985
		US	4060612 A	29-11-1977
		BE	848612 A	23-05-1977
		JP	1356666 C	13-01-1987
		JP	52065254 A	30-05-1977
		JP	61020536 B	22-05-1986
		US	4060641 A	29-11-1977
		JP	1256359 C	12-03-1985
		JP	52085156 A	15-07-1977
		JP	59031492 B	02-08-1984
		US	4100299 A	11-07-1978

DE 1269134 B		NONE		

US 3947434 A	30-03-1976	US	3919238 A	11-11-1975
		AU	475718 B	02-09-1976
		AU	6523674 A	04-09-1975
		BE	816003 A	06-12-1974
		DE	2427409 A	09-01-1975
		DK	301974 A,B,	03-02-1975
		FR	2232313 A	03-01-1975
		GB	1398508 A	25-06-1975
		IL	44141 A	31-08-1976
		JP	50019777 A	01-03-1975

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 40 1944

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-12-1998

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3947434	A		NL 7404135	10-12-1974
			SE 391925	07-03-1977
			SE 7405652	09-12-1974
			ZA 7400683	24-09-1975

US 4751302	A	14-06-1988	JP 58159471	21-09-1983
			JP 58159472	21-09-1983
			JP 58159473	21-09-1983
			JP 58159474	21-09-1983
			JP 58159475	21-09-1983
			JP 58159476	21-09-1983
			JP 58159477	21-09-1983
			EP 0090972	12-10-1983
			US 4533731	06-08-1985

GB 924961	A		BE 588558	
			CH 361005	
			CH 395126	
			CH 436259	
			DE 1238485	
			FR 558	
			FR 84256	05-05-1965
			FR 1421206	09-03-1966
			GB 824853	
			GB 921978	
			LU 38374	
NL 129619				
NL 249341				

US 3312696	A	04-04-1967	NONE	

DE 2624261	A	16-12-1976	FR 2313042	31-12-1976
			FR 2349332	25-11-1977
			BE 842453	01-10-1976
			CH 597192	31-03-1978
			GB 1513092	07-06-1978

US 2870151	A		NONE	

DE 965813	C		NONE	
